

ERRATA:
SHIMURA CURVES OF GENUS AT MOST TWO

JOHN VOIGHT

This note gives errata for the article *Shimura curves of genus at most two* [3]. The author would like to thank Steve Donnelly, Virgile Ducet, James Rickards, and Eloi Torrents Juste for corrections.

ERRATA

- (1) In the script to count, a human error caused the classical modular curve $X_0(11)$ to be counted as genus 0. Moreover, the code which removed duplicates had bugs in redundancies. The code is now available [4]. The corrected counts are 257, 338, 273 of genus 0, 1, 2 up to equivalence, and 326, 467, 416 of genus 0, 1, 2 fixing a unique field in each isomorphism class.
- (2) Section 2, second paragraph: this is not a bijection, but a two-to-one map—an embedding and its conjugate give the same *embedded* finite subgroup.
- (3) Lemma 2.1: The factor 2 from the previous mistake cancels the 2 in the denominator, so the corrected formula is:

$$e_q = \frac{1}{h(F)} \sum_{\substack{R \subset K_q \\ w(R)=2q}} \frac{h(R)}{Q(R)} \prod_{\mathfrak{p}} m(R_{\mathfrak{p}}, \mathcal{O}_{\mathfrak{p}}).$$

(The formula that was implemented included both of these mistakes, the effect of which cancelled out.)

- (4) Lemma 2.1: should be $Q(R) = [N_{K_q/F}(R^*) : \mathbb{Z}_F^{*2}]$.
- (5) Corollary 2.4: Same as the previous error: we have instead

$$e_q = \frac{1}{h(F)} \prod_{\mathfrak{p}|\mathfrak{D}} \left(1 - \left(\frac{K_q}{\mathfrak{p}} \right) \right) \sum_{\substack{R \subset K_q \\ w(R)=2q}} \frac{h(R)}{Q(R)}.$$

- (6) Lemma 2.5 is incorrect as stated. The detailed correction is given in the next section.

EMBEDDING NUMBERS

Lemma 2.5 is incorrect as stated. This mistake does not affect any other result in the paper; the list of curves is still complete and all curves have the correct signature (these signatures were independently verified, as in §5).

We give a complete and corrected statement and proof below. We retain the notation from §2. In particular, let $R_{\mathfrak{p}} = \mathbb{Z}_{F,\mathfrak{p}}[\gamma_{\mathfrak{p}}]$ and let π be a uniformizer at \mathfrak{p} , and let $f_{\mathfrak{p}}(x) = x^2 - t_{\mathfrak{p}}x + n_{\mathfrak{p}}$ denote the minimal polynomial of $\gamma_{\mathfrak{p}}$. Let $d_{\mathfrak{p}} = t_{\mathfrak{p}}^2 - 4n_{\mathfrak{p}}$ and let $k(\mathfrak{p})$ denote the residue class field of \mathfrak{p} .

We will use the following proposition in the proof; see Hijikata [1, §2] and Vignéras [2, §III.3].

Proposition 2.3 (Hijikata [1, Theorem 2.3]).

(c) Suppose $\mathfrak{p} \mid \mathfrak{N}$. Let $e = \text{ord}_{\mathfrak{p}}(\mathfrak{N})$ and for $s \geq e$ let

$$E(s) = \{x \in \mathbb{Z}_F/\mathfrak{p}^s : f_{\mathfrak{p}}(x) \equiv 0 \pmod{\mathfrak{p}^s}\}.$$

If $\text{ord}_{\mathfrak{p}}(d_{\mathfrak{p}}) = 0$ then

$$m(R_{\mathfrak{p}}, \mathcal{O}_{\mathfrak{p}}) = \#E(e).$$

Otherwise,

$$m(R_{\mathfrak{p}}, \mathcal{O}_{\mathfrak{p}}) = \#E(e) + \#\text{img}(E(e+1) \rightarrow \mathbb{Z}_F/\mathfrak{p}^e).$$

The corrected lemma is then as follows.

Lemma 2.5. Let \mathfrak{p} be an odd prime. Suppose $e = \text{ord}_{\mathfrak{p}}(\mathfrak{N}) \geq 1$, let $r = \text{ord}_{\mathfrak{p}}(d_{\mathfrak{p}})$, and let $\kappa = \#k(\mathfrak{p})$.

• If $r = 0$, then

$$m(R_{\mathfrak{p}}, \mathcal{O}_{\mathfrak{p}}) = 1 + \left(\frac{K_q}{\mathfrak{p}}\right).$$

• If $e < r$, then

$$m(R_{\mathfrak{p}}, \mathcal{O}_{\mathfrak{p}}) = \begin{cases} 2\kappa^{(e-1)/2}, & \text{if } e \text{ is odd;} \\ \kappa^{e/2-1}(\kappa+1), & \text{if } e \text{ is even.} \end{cases}$$

• If $e = r$, then

$$m(R_{\mathfrak{p}}, \mathcal{O}_{\mathfrak{p}}) = \begin{cases} \kappa^{(r-1)/2}, & \text{if } r \text{ is odd;} \\ \kappa^{r/2} + \kappa^{r/2-1} \left(1 + \left(\frac{K_q}{\mathfrak{p}}\right)\right), & \text{if } r \text{ is even.} \end{cases}$$

• If $e > r > 0$, then

$$m(R_{\mathfrak{p}}, \mathcal{O}_{\mathfrak{p}}) = \begin{cases} 0, & \text{if } r \text{ is odd;} \\ \kappa^{r/2-1}(\kappa+1) \left(1 + \left(\frac{K_q}{\mathfrak{p}}\right)\right), & \text{if } r \text{ is even.} \end{cases}$$

Proof. Since \mathfrak{p} is odd, without loss of generality we may assume that $\text{trd}(\gamma_{\mathfrak{p}}) = 0$, and hence $E(s)$ is in bijection with

$$E(s) = \{x \in \mathbb{Z}_F/\mathfrak{p}^s : x^2 \equiv d_{\mathfrak{p}} \pmod{\mathfrak{p}^s}\}.$$

First suppose $r = 0$. By Proposition 2.3(c), we have $m = m(R_{\mathfrak{p}}, \mathcal{O}_{\mathfrak{p}}) = \#E(e)$, and by Hensel's lemma we see that $\#E(e) = 0$ or 2 according as $d_{\mathfrak{p}}$ is a square or not in $\mathbb{Z}_{F,\mathfrak{p}}$. In all other cases, we have the second case of Proposition 2.3(c).

Now suppose that $e < r$. The solutions to the equation $x^2 \equiv 0 \pmod{\mathfrak{p}^s}$ are those with $x \equiv 0 \pmod{\mathfrak{p}^{\lceil s/2 \rceil}}$. Thus $\#E(e) = \kappa^{e-\lceil e/2 \rceil} = \kappa^{\lfloor e/2 \rfloor}$ and we see that $\#\text{img}(E(e+1) \rightarrow R/\mathfrak{p}^e) = \kappa^{e-\lceil (e+1)/2 \rceil}$, so $m = 2\kappa^{(e-1)/2}$ if e is odd and $m = \kappa^{e/2} + \kappa^{e/2-1} = \kappa^{e/2-1}(\kappa+1)$ if e is even.

If $e = r$, then again $\#E(e) = \kappa^{\lfloor e/2 \rfloor}$. Now to count the second contributing set, we must solve $x^2 \equiv d_{\mathfrak{p}} \pmod{\mathfrak{p}^{e+1}}$. If $e = r$ is odd then this congruence has no solution. If instead e is even then we must solve $y^2 = (x/\pi^{r/2})^2 \equiv d_{\mathfrak{p}}/\pi^r \pmod{\mathfrak{p}}$ where π is a uniformizer at \mathfrak{p} . This latter congruence has zero or two solutions according as $d_{\mathfrak{p}}$ is a square, and given such a solution y we have the

solutions $x \equiv y \pmod{\pi^{r/2+1}}$ to the original congruence, and hence there are 0 or $2\kappa^{r-(r/2+1)} = 2\kappa^{r/2-1}$ solutions, as claimed.

Finally, suppose $e > r > 0$. If r is odd, there are no solutions to $x^2 \equiv d_{\mathfrak{p}} \pmod{\mathfrak{p}^e}$. If r is even, there are no solutions if $d_{\mathfrak{p}}$ is not a square and otherwise the solutions are $x \equiv y \pmod{\mathfrak{p}^{e-r/2}}$ as above so they total $2\kappa^{r/2} + 2\kappa^{r/2-1} = 2\kappa^{r/2-1}(\kappa + 1)$. \square

REFERENCES

- [1] Hiroaki Hijikata, *Explicit formula of the traces of Hecke operators for $\Gamma_0(N)$* , J. Math. Soc. Japan **26** (1974), no. 1, 56–82.
- [2] Marie-France Vignéras, *Arithmétique des algèbres de quaternions*, Lecture Notes in Math., vol. 800, Springer, Berlin, 1980.
- [3] John Voight, Shimura curves of genus at most two, *Math. Comp.* **78** (2009), 1155–1172.
- [4] John Voight, shim-enum package, <https://github.com/jvoight/shim-enum>, 2025.